

PROSETTA STONE[®] ANALYSIS REPORT

A ROSETTA STONE FOR PATIENT REPORTED OUTCOMES

PROMIS PEDIATRIC FATIGUE AND PEDIATRIC FACIT
FATIGUE

DAVID CELLA, BENJAMIN D. SCHALET, MICHAEL A. KALLEN, JIN-SHEI LAI, KARON F. COOK, JOSHUA
RUTSOHN & SEUNG W. CHOI

DEPARTMENT OF MEDICAL SOCIAL SCIENCES
FEINBERG SCHOOL OF MEDICINE
NORTHWESTERN UNIVERSITY

This research was supported by an NIH/National Cancer Institute grant PROSETTA STONE (1RC4CA157236-01, PI: David Cella). Authors acknowledge careful reviews, comments, and suggestions from Drs. Robert Brennan, Lawrence Hedges, Won-Chan Lee, and Nan Rothrock.

Table of Contents

| | |
|--|----|
| 1. Introduction | 3 |
| 2. The PRO Rosetta Stone Project | 3 |
| 2.1. Patient-Reported Outcomes Measurement Information System (PROMIS) | 4 |
| 2.2. The NIH Toolbox for Assessment of Neurological and Behavioral Function (NIH Toolbox)..... | 5 |
| 2.3. Quality of Life Outcomes in Neurological Disorders (Neuro-QoL) | 5 |
| 3. Legacy Instruments | 6 |
| 3.3 Pediatric Functional Assessment of Chronic Illness Therapy – Fatigue (pedsFACIT-Fatigue) | 6 |
| 4. Linking Methods..... | 6 |
| 4.1. IRT Linking | 7 |
| 4.2. Equipercentile Linking..... | 8 |
| 4.3. Linking Assumptions | 9 |
| 5. Linking Results | 9 |
| 5.5 PROMIS Pediatric Fatigue and Pediatric FACIT Fatigue..... | 11 |
| 5.5.1 Raw Summed Score Distribution..... | 11 |
| 5.5.2 Classical Item Analysis | 12 |
| 5.5.3 Confirmatory Factor Analysis (CFA)..... | 12 |
| 5.5.4 Item Response Theory (IRT) Linking | 13 |
| 5.5.5 Raw Score to T-Score Conversion using Linked IRT Parameters | 15 |
| 5.5.6 Equipercentile Linking..... | 15 |
| 5.5.7 Summary and Discussion..... | 16 |
| 6. Appendix Table 13: Raw Score to T-Score Conversion Table (IRT Fixed Parameter Calibration Linking) for Pediatric FACIT Fatigue and PROMIS Pediatric Fatigue (PROsetta Stone Wave 2 Study) – RECOMMENDED | 19 |
| 7. Appendix Table 14: Direct (Raw to Scale) Equipercentile Crosswalk Table – From Pediatric FACIT Fatigue to PROMIS Pediatric Fatigue – Table 13 is recommended..... | 21 |
| 8. Appendix Table 15: Indirect (Raw to Raw to Scale) Equipercentile Crosswalk Table – Pediatric FACIT Fatigue to PROMIS Pediatric Fatigue – Table 13 is recommended..... | 23 |

PRO Rosetta Stone (*PROsetta Stone*®) Analysis

1. Introduction

A common problem when using a variety of patient-reported outcome measures (PROs) for diverse populations and subgroups is establishing the comparability of scales or units on which the outcomes are reported. The lack of comparability in metrics (e.g., raw summed scores vs. scaled scores) among different PROs poses practical challenges in measuring and comparing effects across different studies. Linking refers to establishing a relationship between scores on two different measures that are not necessarily designed to have the same content or target population. When tests are built in such a way that they differ in content or difficulty, linking must be conducted in order to establish a relationship between the test scores. One technique, commonly referred to as equating, involves the process of converting the system of units of one measure to that of another. This process of deriving equivalent scores has been used successfully in educational assessment to compare test scores obtained from parallel or alternate forms that measure the same characteristic with equal precision. Extending the technique further, comparable scores are sometimes derived for measures of different but related characteristics. The process of establishing comparable scores generally has little effect on the magnitude of association between the measures. Comparability may not signify interchangeability unless the association between the measures approaches unit reliability. Equating, the strongest form of linking, can be established only when two tests 1) measure the same content/construct, 2) target very similar populations, 3) are administered under similar conditions such that the constructs measured are not differentially affected, 4) share common measurement goals and 5) are equally reliable. When test forms are created to be similar in content and difficulty, equating adjusts for differences in difficulty. Test forms are then considered to be essentially the same, so scores on the two forms can be used interchangeably after equating has adjusted for differences in difficulty. For tests with lesser degrees of similarity, only weaker forms of linking are meaningful, such as calibration, concordance, projection, or moderation.

2. The PRO Rosetta Stone Project

The primary aim of the PRO Rosetta Stone (*PROsetta Stone*®) project (1RC4CA157236-01, PI: David Cella) is to develop and apply methods to link the Patient-Reported Outcomes Measurement Information System (PROMIS) measures with other related “legacy” instruments in order to expand the range of PRO assessment options within a common, standardized metric. The project identifies and applies appropriate linking methods that allow scores on a range of legacy PRO instruments to be expressed as standardized T-score metrics linked to the PROMIS metric. This report (Volume 3) encompasses 8 linking studies based on available pediatric PRO data from NIH Toolbox, Neuro-QoL, and *PROsetta Stone* Wave 3. The *PROsetta Stone* Report Volume 1 included linking results primarily from PROMIS Wave 1, as well as links based on NIH Toolbox and Neuro-QoL data. Volume 2 included linking studies based on data that were primarily from *PROsetta Stone* Waves 1 and 2.

2.1. Patient-Reported Outcomes Measurement Information System (PROMIS)

In 2004, the NIH initiated the PROMIS¹ cooperative group under the NIH Roadmap² effort to re-engineer the clinical research enterprise. The aim of PROMIS is to revolutionize and standardize how PRO tools are selected and employed in clinical research. To accomplish this, a publicly-available system was developed to allow clinical researchers access to a common repository of items and state-of-the-science computer-based methods for administering the PROMIS measures. The PROMIS measures include item banks across a wide range of domains that comprise physical, mental, and social health for adults and children, with 12-124 items per bank. Initial concepts measured include emotional distress (anger, anxiety, and depression), physical function, fatigue, pain (quality, behavior, and interference), social function, sleep disturbance, and sleep-related impairment. The banks can be used to administer computerized adaptive tests (CAT) or fixed-length forms in these domains. We have also developed 4-item to 20-item short forms for each domain, and a 10-item Global Health Scale that includes global ratings of five broad PROMIS domains and general health perceptions. As described in a full issue of *Medical Care* (Cella et al., 2007), the PROMIS items, banks, and short forms were developed using a standardized, rigorous methodology that began with constructing a consensus-based PROMIS domain framework.

All PROMIS banks have been calibrated according to Samejima's (1969) graded response model and are based on large data collections including both general and clinical samples. All PROMIS banks are re-scaled (mean=50 and SD=10) using scale-setting subsamples matching the marginal distributions of gender, age, race, and education in the 2000 US census. The PROMIS Wave I calibration data included (a) a small number of full-bank testing cases (approximately 1,000 per bank) from a general population taking one full bank and (b) a larger number of block-administration cases (n= ~14,000) from both general and clinical populations taking a collection of blocks representing all banks, with seven items administered from each bank. The full-bank testing samples were randomly assigned to one of seven different forms. Each form was composed of one or more PROMIS domains (with an exception of Physical Function, where the bank was split over two forms) and one or more legacy measures of the same or related domains.

The PROMIS Wave I data collection design included a number of widely accepted "legacy" measures. The legacy measures used for validation evidence included Buss-Perry Aggression Questionnaire (BPAQ), Center for Epidemiological Studies Depression Scale (CES-D), Mood and Anxiety Symptom Questionnaire (MASQ), Functional Assessment of Chronic Illness Therapy-Fatigue (FACIT-F), Brief Pain Inventory (BPI), and SF-36. In addition to PROMIS-legacy measure pairings for validity assessment (e.g., PROMIS Depression and CES-D), the PROMIS Wave I data allowed for the potential to link over a dozen pairs of measures/subscales. Furthermore, included within each of the PROMIS banks were items from many other existing measures. Depending on the nature and strength of relationship between the measures, various linking procedures can be used to allow for cross-walking of scores. (Note that most of the linking reports based on the PROMIS Wave 1 dataset are included in Volume 1.)

¹ www.nihpromis.org

² www.nihroadmap.nih.gov

2.2. The NIH Toolbox for Assessment of Neurological and Behavioral Function (NIH Toolbox)

Developed in 2006 with the NIH Blueprint funding for Neuroscience Research, four domains of assessment central to neurological and behavioral function were created to measure cognition, sensation, motor functioning, and emotional health. The NIH Toolbox for Assessment of Neurological and Behavioral Function³ provides investigators with brief, yet comprehensive measurement tools for assessment of cognitive function, emotional health, sensory, and motor function. It provides an innovative approach to measurement that is responsive to the needs of researchers in a variety of settings, with a particular emphasis on measuring outcomes in clinical trials and functional status in large cohort studies (e.g., epidemiological studies and longitudinal studies). Included as subdomains of emotional health were negative affect, psychological well-being, stress and self-efficacy, and social relationships. Three PROMIS emotional distress item banks (Anger, Anxiety, and Depression) were used as measures of negative affect. Additionally, existing “legacy” measures, e.g., Patient Health Questionnaire (PHQ-9) and Center for Epidemiological Studies Depression Scale (CES-D), were flagged as potential candidates for the NIH Toolbox battery because of their history, visibility, and research legacy. Among these legacy measures, we focused on those that were available without proprietary restrictions for research applications. In most cases, these measures had been developed using classical test theory.

2.3. Quality of Life Outcomes in Neurological Disorders (Neuro-QoL)

The National Institute of Neurological Disorders and Stroke sponsored a multi-site project to develop clinically relevant and psychometrically robust Quality of Life (QOL) assessment tools for adults and children with neurological disorders. The primary goal of this effort, known as Neuro-QoL⁴, was to enable clinical researchers to compare the QOL impact of different interventions within and across various conditions. This resulted in 13 adult QOL item banks (Anxiety, Depression, Fatigue, Upper Extremity Function - Fine Motor, Lower Extremity Function - Mobility, Applied Cognition - General Concerns, Applied Cognition - Executive Function, Emotional and Behavioral Dyscontrol, Positive Affect and Well-Being, Sleep Disturbance, Ability to Participate in Social Roles and Activities, Satisfaction with Social Roles and Activities, and Stigma), eight pediatric item banks (Anger, Anxiety, Depression, Fatigue, Pain, Applied Cognition - General Concerns, Social Relations - Interaction with Peers, and Stigma) and two additional pediatric physical function scales (Lower Extremity Function - Mobility, and Upper Extremity Function - Fine Motor, ADL).

³ www.nihtoolbox.org

⁴ www.neuroqol.org

3. Legacy Instruments

The following instruments are widely accepted “legacy” measures that have now been linked to PROMIS instruments. Some of these legacy measures were used as part of the initial validation work for PROMIS and NIH Toolbox. In some cases, instruments were administered as part of the PROsetta Stone project for specific linking purposes. Data were collected on reference measures (e.g., PROMIS Depression) from a minimum of 400 respondents (for stable item parameter estimation), along with responses to at least one other conceptually similar scale or bank to be linked to the reference measure. (See Table 5.1).

3.3 Pediatric Functional Assessment of Chronic Illness Therapy - Fatigue (pedsFACIT-Fatigue)

The Pediatric Functional Assessment of Chronic Illness Therapy-Fatigue (pedsFACIT-F) is a 13-item self-report measure of pediatric cancer-related fatigue and is part of the Functional Assessment of Chronic Illness Therapy (FACIT) measurement system. pedsFACIT-F was derived from the 51-item pediatric fatigue item bank (pedsFIB). Each item is rated on a 0 to 4 scale: "None of the time", "A little bit of the time", "Some of the time", "Most of the time", "All of the time". It measures fatigue in the past seven days. pedsFACIT-F deals with the issue of different literacy levels as children age, as well as the differing perceptions of the impact of fatigue over time. Because of the unique aspects of assessing cancer-related fatigue among children, simply modifying the item wording to make the language more developmentally appropriate may not be sufficient for providing clinically useful information in monitoring fatigue over time. pedsFACIT-F has stable measurement properties across age, sex, and cancer types. It can be used in clinical research. Scores on the pedsFACIT-F discriminate between patients with and without anemia and among patients with different functional status. As with all FACIT questionnaires, a high score is good. Therefore, a score of “0” indicates a severely symptomatic patient, and the highest possible score (varies per scale and subscale) indicates an asymptomatic patient.

4. Linking Methods

PROMIS full-bank administration allows for single-group linking. This linking method is used when two or more measures are administered to the same group of people. For example, two PROMIS banks (Anxiety and Depression) and three legacy measures (MASQ, CES-D, and SF-36 MH) were administered to a sample of 925 people, with the order of measures presented randomized so as to minimize potential order effects. The original purpose of the PROMIS full-bank administration study was to establish initial validity evidence (e.g., validity coefficients), not to establish linking relationships. Thus, initial analyses of the full-bank administration sample revealed several potential score-linking issues: (a) some measures had severely skewed score distributions; (b) the sample size for some administered measures was relatively small. These score-linking issues can be limiting factors when determining an appropriate linking method (e.g., what method options are available or whether linking can even be conducted). Another potential linking issue is related to how the non-PROMIS measures are scored and reported. For example, all SF-36 subscales are scored using a proprietary scoring algorithm and reported

as normed scores (0 to 100). Others are scored and reported using simple raw summed scores. All PROMIS measures are scored using the final re-centered item response theory (IRT) item parameters and transformed to the T-score metric (mean=50, SD=10).

PROMIS's T-score distributions are standardized so that a score of 50 represents the average (mean) for the US general population and the standard deviation around that mean is 10 points. A high PROMIS score always represents more of the concept being measured. Thus, a person who has a T-score of 60 is one standard deviation higher than the general population for the concept being measured. It therefore follows that, for condition symptoms and negatively-framed or oriented concepts like pain, fatigue, and anxiety, a score of 60 is one standard deviation worse than average; while for functional scores and positively-framed or oriented concepts like physical and social function, a score of 60 is one standard deviation better than average.

In order to apply linking methods consistently across different studies, linking/concordance relationships were established based on the raw summed score metric of the measures. Furthermore, the direction of linking relationships established was from legacy to PROMIS measure. That is, each raw summed score on a given legacy instrument was mapped to a T-score on the corresponding PROMIS instrument/bank. Finally, the raw summed score for each legacy instrument was constructed so that higher scores would represent higher levels of the construct being measured (to be consistent with the PROMIS approach). When legacy measures were scaled in the opposite direction, we reversed the direction of the legacy measure in order for the correlation between legacy and PROMIS measures to be positive and thereby facilitate concurrent calibration. As a result, some or all item response scores for some legacy instruments needed to be reverse-coded.

4.1. IRT Linking

One of the objectives of the current linking analyses is to determine whether the non-PROMIS measures can be added to their respective PROMIS item banks without significantly altering the underlying trait being measured. The rationale is twofold: (1) the augmented PROMIS item banks might provide more robust coverage, both in terms of content and difficulty; and (2) calibrating the non-PROMIS measures on the corresponding PROMIS item bank scale might facilitate subsequent linking analyses. At least two IRT linking approaches are applicable under the current study design: (1) linking separate calibrations through the Stocking-Lord method and (2) fixed-parameter calibration.

Linking separate calibrations might involve the following steps under the current setting.

- First, simultaneously calibrate the combined item set (e.g., PROMIS Depression bank and CES-D).
- Second, estimate linear transformation coefficients (additive and multiplicative constants) using the item parameters for the PROMIS bank items as anchor items.
- Third, transform the metric for the non-PROMIS items to the PROMIS metric.

The second approach, fixed-parameter calibration, involves fixing the PROMIS item parameters at their final bank values and calibrating only non-PROMIS items in order that the non-PROMIS item parameters may be placed on the same metric as the PROMIS items; that is, the focus is

on placing the parameters of non-PROMIS items on the PROMIS metric. Updating the PROMIS item parameters is not desired, because the larger PROsetta-wide linking exercise is built on the stability of these final PROMIS calibrations. Note that IRT linking would be necessary when the ability level of the full-bank testing sample is different from that of the PROMIS scale-setting sample. If it is assumed that the two samples are from the same population, linking is not necessary and calibration of the items (either separately or simultaneously) will result in item parameter estimates that are on the same scale metric without any further scale linking. Even though the full-bank testing sample was a subset of the full PROMIS calibration sample, it is still possible that the two samples are somewhat disparate due to some non-random component of the selection process. Moreover, there is some evidence that linking can improve the accuracy of parameter estimation even when linking is not fully necessary (e.g., two samples are from the same population having the same or similar ability levels). Thus, conducting IRT linking would be worthwhile, with potential score accuracy benefits gained.

Once the non-PROMIS items are calibrated on the corresponding PROMIS item bank metric, the augmented item bank can be used for standard computation of IRT scaled scores from any subset of the items, including computerized adaptive testing (CAT) and creating short forms. The non-PROMIS items will be treated the same as the existing PROMIS items. Again, the above options are feasible only when the dimensionality of the bank is not altered significantly (i.e., where a unidimensional IRT model remains suitable for the aggregate set of items). Thus, prior to conducting IRT linking, it is important to assess the dimensionality of the involved measures based on separate and combined PROMIS and non-PROMIS measures. Various dimensionality assessment tools can be used, including confirmatory factor analysis, disattenuated correlations, and essential unidimensionality.

4.2. Equipercentile Linking

The IRT linking procedures described above are permissible only if the traits being measured are not significantly altered by aggregating items from multiple measures. One potential issue might be the creation of multidimensionality as a result of aggregating items measuring different traits. For two scales that measure distinct but highly related traits, predicting scores on one scale from those of the other has been used frequently. Concordance tables between PROMIS and non-PROMIS measures can be constructed using equipercentile equating (Lord, 1982; Kolen & Brennan, 2004) when there is insufficient empirical evidence that the instruments measure the same construct. An equipercentile method estimates a nonlinear linking relationship using percentile rank distributions of the two linking measures. The equipercentile linking method can be used in conjunction with a presmoothing method such as the loglinear model (Hanson, Zeng, & Colton, 1994). The frequency distributions are first smoothed using the loglinear model and then equipercentile linking is conducted based on the smoothed frequency distributions of the two measures. Smoothing can also be done at the backend on equipercentile equivalents and is called postsmoothing (Brennan, 2004; Kolen & Brennan, 2004). The cubic-spline smoothing algorithm (Reinsch, 1967) is used in the LEGS program employed in PROsetta analyses (Brennan, 2004). Smoothing is intended to reduce sampling error involved in the linking process. A successful linking procedure will provide a conversion (crosswalk) table, in which, for example, raw summed scores on the PHQ-9 measure are transformed to the T-score equivalents of the PROMIS Depression measure.

In the current context, equipercentile crosswalk tables can be generated using two different approaches. First is a direct linking approach where each raw summed score on a non-PROMIS measure is mapped directly to a PROMIS T-score. That is, raw summed scores on the non-PROMIS instrument and IRT scaled scores on the PROMIS (reference) instrument are linked directly, although raw summed scores and IRT scaled scores have distinct properties (e.g., discrete vs. continuous). This approach might be appropriate when the reference instrument is either an item bank or composed of a large number of items and so various subsets (static or dynamic) are likely to be used but not the full bank in its entirety (e.g., the PROMIS Physical Function bank with 124 items). Second is an indirect approach where raw summed scores on the non-PROMIS instrument are mapped to raw summed scores on the PROMIS instrument, and then the resulting raw summed score equivalents are mapped to corresponding scaled scores based on a raw-to-scale score conversion table. Because the raw summed score equivalents may take fractional values, such a conversion table will need to be interpolated using statistical procedures (e.g., cubic spline).

Finally, when samples are small or inadequate for a specific method, random sampling error becomes a major concern (Kolen & Brennan, 2004). That is, substantially different linking relationships might be obtained if linking is conducted repeatedly over different samples. This type of random sampling error can be measured by the standard error of equating (SEE), which can be operationalized as the standard deviation of equated scores for a given raw summed score over replications (Lord, 1982).

4.3. Linking Assumptions

In Section 5 of this PROsetta Stone report, we present the results of a large number of linking studies using secondary data sets. In most cases, we have applied all three linking methods described in sections 4.1 and 4.2. Our purpose is to provide the maximum amount of useful information. However, the suitability of these methods depends upon the meeting of various linking assumptions. These assumptions require that the two instruments to be linked measure the same construct, show a high correlation, and are relatively invariant in subpopulation differences (Dorans, 2007). The degree to which these assumptions are met varies across linking studies. Given that different researchers may interpret these requirements differently, we have taken a liberal approach for inclusion of linkages in this book. Nevertheless, we recommend that researchers diagnostically review the classical psychometrics and CFA results in light of these assumptions prior to any application of the cross-walk charts or legacy parameters to their own data. Having investigated a large number of possible links between PROMIS measures and legacy measures, we did apply a few minimal exclusion rules before linking. For example, we generally did not proceed with planned linking when the raw score correlation between two instruments was less than .70.

5. Linking Results

Table 5.1 lists the linking analyses included in this report, which have been conducted based on samples from a PROsetta Stone study (see Section 2 for more details). In most cases, PROMIS instruments were used as the reference (i.e., scores on non-PROMIS instruments are expressed on the PROMIS score metric).

Table 5.1. Linking by Reference Instrument

| Section | PROMIS Instrument | Instrument to Link | Study |
|----------------|--------------------------|---------------------------|-----------------------|
| 5.5 | PROMIS Pediatric Fatigue | Pediatric FACIT Fatigue | PROsetta Stone Wave 3 |

5.5 PROMIS Pediatric Fatigue and Pediatric FACIT Fatigue

In this section we provide a summary of the procedures employed to establish a crosswalk between two measures of fatigue, namely the PROMIS Pediatric Fatigue item bank (15 items) and Peds FACIT Fatigue (13 items). Both instruments were scaled such that higher scores represent higher levels of fatigue. We created raw summed scores for each of the measures separately and then for them combined. Summing of item scores assumes that all items have positive correlations with the total as examined in the section on Classical Item Analysis. Our sample consisted of 507 participants (N = 505 for participants with complete responses).

5.5.1 Raw Summed Score Distribution

The maximum possible raw summed scores were 75 for PROMIS Pediatric Fatigue and 65 for Peds FACIT-F. Figure 5.5.1 and Figure 5.5.2 graphically display the raw summed score distributions of the two measures. Figure 5.5.3 shows the distribution for them combined. Figure 5.5.4 is a scatter plot matrix showing the relationship of each pair of raw summed scores. Pearson correlations are shown above the diagonal. The correlation between PROMIS Pediatric Fatigue and Peds FACIT-F was 0.86. The disattenuated (corrected for unreliabilities) correlation between PROMIS Pediatric Fatigue and Peds FACIT-F was 0.9. The correlations between the combined score and the measures were 0.97 and 0.95 for PROMIS Pediatric Fatigue and Peds FACIT-F, respectively.




Figure 5.5.1: Raw Summed Score Distribution - PROMIS Peds Fatigue




Figure 5.5.2: Raw Summed Score Distribution - Peds FACIT-F




Figure 5.5.3: Raw Summed Score Distribution – Combined




Figure 5.5.4: Scatter Plot Matrix of Raw Summed Scores

5.5.2 Classical Item Analysis

We conducted classical item analyses on the two measures separately and on them combined. Table 5.5.1 summarizes the results. For PROMIS Pediatric Fatigue, Cronbach’s alpha internal consistency reliability estimate was 0.96 and adjusted (corrected for overlap) item-total correlations ranged from 0.576 to 0.833. For Peds FACIT-F, alpha was 0.941 and adjusted item-total correlations ranged from 0.506 to 0.818. For the 28 items, alpha was 0.972 and adjusted item-total correlations ranged from 0.474 to 0.825.

Table 5.5.1: Classical Item Analysis

| | No. Items | Cronbach's Alpha Internal Consistency Reliability Estimate | Adjusted (corrected for overlap) Item-total Correlation | | |
|--------------------------|-----------|--|---|-------|---------|
| | | | Minimum | Mean | Maximum |
| PROMIS Pediatric Fatigue | 15 | 0.960 | 0.576 | 0.770 | 0.833 |
| Peds FACIT-F | 13 | 0.941 | 0.506 | 0.718 | 0.818 |
| Combined | 28 | 0.972 | 0.474 | 0.735 | 0.825 |

5.5.3 Confirmatory Factor Analysis (CFA)

To assess the dimensionality of the measures, a categorical confirmatory factor analysis (CFA) was carried out using the WLSMV estimator of Mplus on a subset of cases without missing responses. A single factor model (based on polychoric correlations) was run on each of the two measures separately and on the combined. Table 5.5.2 summarizes the model fit statistics. For PROMIS Pediatric Fatigue, the fit statistics were as follows: CFI = 0.972, TLI = 0.967, and RMSEA = 0.118. For Peds

FACIT-F, CFI = 0.949, TLI = 0.939, and RMSEA = 0.161. For the 28 items, CFI = 0.951, TLI = 0.948, and RMSEA = 0.102. The main interest of the current analysis is whether the combined measure is essentially unidimensional.

Table 5.5.2: CFA Fit Statistics

| | No. Items | n | CFI | TLI | RMSEA |
|--------------------------|-----------|-----|-------|-------|-------|
| PROMIS Pediatric Fatigue | 15 | 507 | 0.972 | 0.967 | 0.118 |
| PedsFACIT F | 13 | 507 | 0.949 | 0.939 | 0.161 |
| Combined | 28 | 507 | 0.951 | 0.948 | 0.102 |

5.5.4 Item Response Theory (IRT) Linking

We conducted concurrent calibration on the combined set of 28 items according to the graded response model. The calibration was run using `MULTILOG` and two different approaches as described previously (i.e., IRT linking vs. fixed-parameter calibration). For IRT linking, all 28 items were calibrated freely on the conventional theta metric (mean=0, SD=1). Then the 15 PROMIS Pediatric Fatigue items served as anchor items to transform the item parameter estimates for the Peds FACIT-F items onto the PROMIS Pediatric Fatigue metric. We used four IRT linking methods implemented in `plink` (Weeks, 2010): mean/mean, mean/sigma, Haebara, and Stocking-Lord. The first two methods are based on the mean and standard deviation of item parameter estimates, whereas the latter two are based on the item and test information curves. Table 5.5.3 shows the additive (A) and multiplicative (B) transformation constants derived from the four linking methods. For fixed-parameter calibration, the item parameters for the PROMIS Pediatric Fatigue items were constrained to their final bank values, while the Peds FACIT-F items were calibrated, under the constraints imposed by the anchor items.

Table 5.5.3: IRT Linking Constants

| | A | B |
|---------------|-------|--------|
| Mean/Mean | 2.173 | -0.376 |
| Mean/Sigma | 1.453 | 0.164 |
| Haebara | 1.392 | 0.192 |
| Stocking-Lord | 1.618 | 0.018 |

The item parameter estimates for the Peds FACIT-F items were linked to the PROMIS Pediatric Fatigue metric using the transformation constants shown in Table 5.5.3. The Peds FACIT-F item parameter estimates from the fixed-parameter calibration are considered already on the PROMIS Pediatric Fatigue metric. Based on the transformed and fixed-parameter estimates we derived test characteristic curves (TCC) for Peds FACIT-F as shown in Figure 5.5.5. Using the fixed-parameter calibration as a basis we then examined the difference with each of the TCCs from the four linking methods. Figure 5.5.6 displays the differences on the vertical axis.




Figure 5.5.5: Test Characteristic Curves (TCC) from Different Linking Methods




Figure 5.5.6: Difference in Test Characteristic Curves (TCC) Comparison with Fixed-Parameter

Table 5.5.4 shows the fixed-parameter calibration item parameter estimates for Peds FACIT-F. The marginal reliability estimate for Peds FACIT-F based on the item parameter estimates was 0.899. The marginal reliability estimates for PROMIS Pediatric Fatigue and the combined set were 0.873 and 0.94, respectively. The slope parameter estimates for Peds FACIT-F ranged from 1.07 to 2.64 with a mean of 2.06. The slope parameter estimates for PROMIS Pediatric Fatigue ranged from 0.91 to 1.9 with a mean of 1.4. We also derived scale information functions based on the fixed-parameter calibration result. Figure 5.5.7 displays the scale information functions for PROMIS Pediatric Fatigue, Peds FACIT-F, and the combined set of 28. We then computed IRT scaled scores for the three measures based on the fixed-parameter calibration result. Figure 5.5.8 is a scatter plot matrix showing the relationships between the measures.

Table 5.5.4: Fixed-Parameter Calibration Item Parameter Estimates for Pediatric FACIT Fatigue

| a | cb1 | cb2 | cb3 | cb4 | NCAT |
|-------|--------|-------|-------|-------|------|
| 1.189 | -1.539 | 0.977 | 2.263 | 3.442 | 5 |
| 1.067 | -0.676 | 1.623 | 2.867 | 4.333 | 5 |
| 1.430 | -1.999 | 0.142 | 1.731 | 2.893 | 5 |
| 2.158 | -0.682 | 0.696 | 1.801 | 2.596 | 5 |
| 2.059 | -0.805 | 0.635 | 1.812 | 3.175 | 5 |
| 2.039 | -0.308 | 0.741 | 1.764 | 2.715 | 5 |
| 2.318 | -0.080 | 0.883 | 1.801 | 2.723 | 5 |
| 2.558 | 0.063 | 0.883 | 1.885 | 2.768 | 5 |
| 1.885 | 0.007 | 1.132 | 2.251 | 3.469 | 5 |
| 2.636 | 0.291 | 1.187 | 2.061 | 2.523 | 5 |
| 2.422 | 0.652 | 1.424 | 2.288 | 3.194 | 5 |
| 2.592 | 0.339 | 1.049 | 1.872 | 2.598 | 5 |
| 2.445 | 0.543 | 1.197 | 1.921 | 2.759 | 5 |




Figure 5.5.7: Comparison of Scale Information Functions




Figure 5.5.8: Comparison of IRT Scaled Scores

5.5.5 Raw Score to T-Score Conversion using Linked IRT Parameters

The IRT model implemented in PROMIS (i.e., the graded response model) uses the pattern of item responses for scoring, not just the sum of individual item scores. However, a crosswalk table mapping each raw summed score point on Peds FACIT-F to a scaled score on PROMIS Pediatric Fatigue can be useful. Based on the Peds FACIT-F item parameters derived from the fixed-parameter calibration, we constructed a score conversion table. The conversion table displayed in Appendix Table 13 can be used to map simple raw summed scores from Peds FACIT-F to T-score values linked to the PROMIS Pediatric Fatigue metric. Each raw summed score point and corresponding PROMIS scaled score are presented along with the standard error associated with the scaled score. The raw summed score is constructed such that for each item, consecutive integers in base 1 are assigned to the ordered response categories.

5.5.6 Equipercentile Linking

We mapped each raw summed score point on Peds FACIT-F to a corresponding scaled score on PROMIS Pediatric Fatigue by identifying scores on PROMIS Pediatric Fatigue that have the same percentile ranks as scores on Peds FACIT-F. Theoretically, the equipercentile linking function is symmetrical for continuous random variables (X and Y). Therefore, the linking function for the values in X to those in Y is the same as that for the values in Y to those in X. However, for discrete variables like raw summed scores the equipercentile linking functions can be slightly different (due to rounding errors and differences in score ranges) and hence may need to be obtained separately. Figure 5.5.9 displays the cumulative distribution functions of the

measures. Figure 5.5.10 shows the equipercentile linking functions based on raw summed scores, from Peds FACIT-F to PROMIS Pediatric Fatigue. When the number of raw summed score points differs substantially, the equipercentile linking functions could deviate from each other noticeably. The problem can be exacerbated when the sample size is small. Appendix Table 14 and Appendix Table 15 show the equipercentile crosswalk tables. The result shown in Appendix Table 14 is based on the direct (raw summed score to scaled score) approach, whereas Appendix Table 15 shows the result based on the indirect (raw summed score to raw summed score equivalent to scaled score equivalent) approach (Refer to Section 4.2 for details). Three separate equipercentile equivalents are presented: one is equipercentile without post smoothing (“Equipercetile Scale Score Equivalents”) and two with different levels of postsmoothing, i.e., “Equipercetile Equivalents with Postsmoothing (Less Smoothing)” and “Equipercetile Equivalents with Postsmoothing (More Smoothing).” Postsmoothing values of 0.3 and 1.0 were used for “Less” and “More,” respectively (Refer to Brennan, 2004 for details).




Figure 5.5.9: Comparison of Cumulative Distribution Functions based on Raw Summed Scores




Figure 5.5.10: Equipercetile Linking Functions

5.5.7 Summary and Discussion

The purpose of linking is to establish the relationship between scores on two measures of closely related traits. The relationship can vary across linking methods and samples employed. In equipercentile linking, the relationship is determined based on the distributions of scores in a given sample. Although IRT-based linking can potentially offer sample-invariant results, they are based on estimates of item parameters, and hence subject to sampling errors. A potential issue with IRT-based linking methods is, however, the violation of model assumptions as a result of combining items from two measures (e.g., unidimensionality and local independence). As displayed in Figure 5.5.10, the relationships derived from various linking methods are consistent, which suggests that a robust linking relationship can be determined based on the given sample.

To further facilitate the comparison of the linking methods, Table 5.5.5 reports four statistics summarizing the current sample in terms of the differences between the PROMIS Pediatric Fatigue T-scores and Peds FACIT-F scores linked to the T-score metric through different methods. In addition to the seven linking methods previously discussed (see Figure 5.5.10), the method labeled “IRT pattern scoring” refers to IRT scoring based on the pattern of item responses instead of raw summed scores. With respect to the correlation between observed and linked T-scores, IRT pattern scoring produced the best result (0.845), followed by IRT raw-scale (0.839). Similar results were found in terms of the standard deviation of differences and root mean squared difference (RMSD). IRT pattern scoring yielded smallest RMSD (6.909), followed by IRT raw-scale (7.024).

Table 5.5.5: Observed vs. Linked T-scores

| Methods | Correlation | Mean Difference | SD Difference | RMSD |
|--------------------------|--------------------|------------------------|----------------------|-------------|
| IRT pattern scoring | 0.845 | -0.735 | 6.877 | 6.909 |
| IRT raw-scale | 0.839 | -0.775 | 6.988 | 7.024 |
| EQP raw-scale SM=0.0 | 0.838 | 0.597 | 7.273 | 7.291 |
| EQP raw-scale SM=0.3 | 0.836 | 0.877 | 7.471 | 7.515 |
| EQP raw-scale SM=1.0 | 0.836 | 0.957 | 7.513 | 7.566 |
| EQP raw-raw-scale SM=0.0 | 0.838 | 0.503 | 7.245 | 7.256 |
| EQP raw-raw-scale SM=0.3 | 0.837 | 0.635 | 7.299 | 7.320 |
| EQP raw-raw-scale SM=1.0 | 0.836 | 0.715 | 7.367 | 7.394 |

To examine the bias and standard error of the linking results, a resampling study was used. In this procedure, small subsets of cases (e.g., 25, 50, and 75) were drawn with replacement from the study sample (N=507) over a large number of replications (i.e., 10,000).

Table 5.5.6 summarizes the mean and standard deviation of differences between the observed and linked T-scores by linking method and sample size. For each replication, the mean difference between the observed and equated PROMIS Pediatric Fatigue T-scores was computed. Then the mean and the standard deviation of the means were computed over replications as bias and empirical standard error, respectively. As the sample size increased (from 25 to 75), the empirical standard error decreased steadily. At a sample size of 75, IRT pattern scoring produced the smallest standard error, 0.734. That is, the means difference between the mean PROMIS Pediatric Fatigue T-score and the mean equated Peds FACIT-F T-score based on a similar sample of 75 cases is expected to be around ± 1.47 (i.e., 2×0.734).

Table 5.5.6: Comparison of Resampling Results

| Methods | Mean (N=25) | SD (N=25) | Mean (N=50) | SD (N=50) | Mean (N=75) | SD (N=75) |
|--------------------------|----------------|--------------|----------------|--------------|----------------|--------------|
| IRT pattern scoring | -0.753 | 1.343 | -0.735 | 0.919 | -0.741 | 0.734 |
| IRT raw-scale | -0.775 | 1.349 | -0.779 | 0.943 | -0.775 | 0.748 |
| EQP raw-scale SM=0.0 | 0.630 | 1.416 | 0.593 | 0.975 | 0.593 | 0.773 |
| EQP raw-scale SM=0.3 | 0.895 | 1.430 | 0.877 | 1.006 | 0.875 | 0.793 |
| EQP raw-scale SM=1.0 | 0.940 | 1.453 | 0.951 | 1.018 | 0.967 | 0.810 |
| EQP raw-raw-scale SM=0.0 | 0.486 | 1.389 | 0.519 | 0.969 | 0.509 | 0.772 |
| EQP raw-raw-scale SM=0.3 | 0.661 | 1.439 | 0.636 | 0.976 | 0.624 | 0.773 |
| EQP raw-raw-scale SM=1.0 | 0.693 | 1.441 | 0.717 | 0.990 | 0.723 | 0.783 |

Examining a number of linking studies in the current project revealed that the two linking methods (IRT and equipercentile) in general produced highly comparable results. Some noticeable discrepancies were observed (albeit rarely) in some extreme score levels where data were sparse. Model-based approaches can provide more robust results than those relying solely on data when data are sparse. The caveat is that the model should fit the data reasonably well. One of the potential advantages of IRT-based linking is that the item parameters on the linking instrument can be expressed on the metric of the reference instrument, and therefore can be combined without significantly altering the underlying trait being measured. As a result, a larger item pool might be available for computerized adaptive testing or various subsets of items can be used in static short forms. Therefore, IRT-based linking (Appendix Table 13) might be preferred when the results are comparable and no apparent violations of assumptions are evident.

6. Appendix Table 13: Raw Score to T-Score Conversion Table (IRT Fixed Parameter Calibration Linking) for Pediatric FACIT Fatigue and PROMIS Pediatric Fatigue (PROsetta Stone Wave 3 Study) – RECOMMENDED

| Peds FACIT Fatigue Actual Score* | Peds FACIT Fatigue Raw Score* | PROMIS Peds Fatigue T-Score | SE |
|---|--------------------------------------|------------------------------------|-----------|
| 52 | 13 | 29.6 | 5.7 |
| 51 | 14 | 33.5 | 5.1 |
| 50 | 15 | 36.7 | 4.6 |
| 49 | 16 | 39.3 | 4.2 |
| 48 | 17 | 41.5 | 3.9 |
| 47 | 18 | 43.4 | 3.6 |
| 46 | 19 | 45.1 | 3.4 |
| 45 | 20 | 46.5 | 3.2 |
| 44 | 21 | 47.9 | 3.1 |
| 43 | 22 | 49.1 | 3.0 |
| 42 | 23 | 50.3 | 2.9 |
| 41 | 24 | 51.3 | 2.8 |
| 40 | 25 | 52.4 | 2.7 |
| 39 | 26 | 53.3 | 2.7 |
| 38 | 27 | 54.2 | 2.6 |
| 37 | 28 | 55.1 | 2.6 |
| 36 | 29 | 56.0 | 2.6 |
| 35 | 30 | 56.8 | 2.6 |
| 34 | 31 | 57.7 | 2.5 |
| 33 | 32 | 58.5 | 2.5 |
| 32 | 33 | 59.3 | 2.5 |
| 31 | 34 | 60.0 | 2.5 |
| 30 | 35 | 60.8 | 2.5 |
| 29 | 36 | 61.6 | 2.5 |
| 28 | 37 | 62.3 | 2.5 |
| 27 | 38 | 63.1 | 2.5 |
| 26 | 39 | 63.8 | 2.5 |
| 25 | 40 | 64.6 | 2.5 |
| 24 | 41 | 65.3 | 2.5 |
| 23 | 42 | 66.1 | 2.5 |
| 22 | 43 | 66.8 | 2.5 |
| 21 | 44 | 67.6 | 2.5 |
| 20 | 45 | 68.3 | 2.5 |
| 19 | 46 | 69.1 | 2.5 |
| 18 | 47 | 69.8 | 2.5 |
| 17 | 48 | 70.6 | 2.5 |
| 16 | 49 | 71.4 | 2.5 |
| 15 | 50 | 72.1 | 2.5 |
| 14 | 51 | 72.9 | 2.5 |
| 13 | 52 | 73.7 | 2.5 |

PROsetta Stone® - PROMIS Pediatric Fatigue and Pediatric FACIT Fatigue

| | | | |
|----|----|------|-----|
| 12 | 53 | 74.5 | 2.6 |
| 11 | 54 | 75.4 | 2.6 |
| 10 | 55 | 76.3 | 2.6 |
| 9 | 56 | 77.2 | 2.7 |
| 8 | 57 | 78.1 | 2.7 |
| 7 | 58 | 79.1 | 2.8 |
| 6 | 59 | 80.1 | 2.9 |
| 5 | 60 | 81.2 | 2.9 |
| 4 | 61 | 82.4 | 3.0 |
| 3 | 62 | 83.5 | 2.9 |
| 2 | 63 | 84.7 | 2.8 |
| 1 | 64 | 85.7 | 2.6 |
| 0 | 65 | 86.6 | 2.3 |

*The scores in the first column conform to the scoring rules of the Peds FACIT Fatigue instrument. The scores in the second column correspond to the scores used in the full PROsetta Stone report.

7. Appendix Table 14: Direct (Raw to Scale) Equipercentile Crosswalk Table – From Pediatric FACIT Fatigue to PROMIS Pediatric Fatigue – Table 13 is recommended

| Peds FACIT Fatigue Actual Score* | Peds FACIT Fatigue Raw Score* | Equipercentile PROMIS Scaled Score Equivalents (No Smoothing) | Equipercentile Equivalents with Postsmoothing (Less Smoothing) | Equipercentile Equivalents with Postsmoothing (More Smoothing) | Standard Error of Equating (SEE) |
|---|--------------------------------------|--|---|---|---|
| 52 | 13 | 28 | 22 | 22 | 0.09 |
| 51 | 14 | 28 | 28 | 29 | 0.09 |
| 50 | 15 | 32 | 32 | 32 | 0.39 |
| 49 | 16 | 36 | 36 | 35 | 0.67 |
| 48 | 17 | 39 | 39 | 39 | 0.58 |
| 47 | 18 | 42 | 41 | 41 | 0.42 |
| 46 | 19 | 43 | 43 | 44 | 0.86 |
| 45 | 20 | 45 | 45 | 46 | 0.72 |
| 44 | 21 | 47 | 47 | 47 | 0.80 |
| 43 | 22 | 50 | 49 | 49 | 0.37 |
| 42 | 23 | 50 | 50 | 50 | 0.36 |
| 41 | 24 | 51 | 51 | 51 | 0.78 |
| 40 | 25 | 52 | 52 | 52 | 0.35 |
| 39 | 26 | 53 | 53 | 53 | 0.58 |
| 38 | 27 | 54 | 54 | 54 | 1.37 |
| 37 | 28 | 56 | 55 | 55 | 0.37 |
| 36 | 29 | 56 | 56 | 56 | 0.33 |
| 35 | 30 | 58 | 57 | 57 | 0.96 |
| 34 | 31 | 58 | 58 | 58 | 0.88 |
| 33 | 32 | 59 | 59 | 59 | 0.42 |
| 32 | 33 | 59 | 59 | 59 | 0.40 |
| 31 | 34 | 60 | 60 | 60 | 0.82 |
| 30 | 35 | 60 | 60 | 60 | 0.24 |
| 29 | 36 | 61 | 61 | 61 | 0.24 |
| 28 | 37 | 61 | 61 | 62 | 0.23 |
| 27 | 38 | 61 | 62 | 62 | 0.22 |
| 26 | 39 | 63 | 63 | 63 | 0.68 |
| 25 | 40 | 64 | 64 | 64 | 0.45 |
| 24 | 41 | 66 | 65 | 65 | 0.61 |
| 23 | 42 | 66 | 66 | 66 | 0.60 |
| 22 | 43 | 67 | 67 | 67 | 0.53 |
| 21 | 44 | 67 | 67 | 67 | 0.50 |
| 20 | 45 | 68 | 68 | 68 | 0.98 |

PROsetta Stone® - PROMIS Pediatric Fatigue and Pediatric FACIT Fatigue

| | | | | | |
|----|----|----|----|----|------|
| 19 | 46 | 68 | 69 | 69 | 0.88 |
| 18 | 47 | 70 | 70 | 70 | 4.24 |
| 17 | 48 | 72 | 71 | 71 | 0.70 |
| 16 | 49 | 72 | 72 | 72 | 0.57 |
| 15 | 50 | 72 | 72 | 72 | 0.49 |
| 14 | 51 | 72 | 73 | 73 | 0.49 |
| 13 | 52 | 73 | 74 | 74 | 2.45 |
| 12 | 53 | 75 | 75 | 75 | 1.37 |
| 11 | 54 | 77 | 77 | 76 | 2.83 |
| 10 | 55 | 78 | 78 | 78 | 2.45 |
| 9 | 56 | 81 | 80 | 79 | 2.00 |
| 8 | 57 | 86 | 81 | 80 | 1.06 |
| 7 | 58 | 86 | 82 | 81 | 1.06 |
| 6 | 59 | 86 | 83 | 82 | 1.06 |
| 5 | 60 | 86 | 85 | 84 | 1.06 |
| 4 | 61 | 86 | 86 | 85 | 1.06 |
| 3 | 62 | 87 | 87 | 86 | 1.06 |
| 2 | 63 | 88 | 88 | 87 | 1.06 |
| 1 | 64 | 89 | 89 | 89 | 1.06 |
| 0 | 65 | 90 | 90 | 90 | 1.06 |

*The scores in the first column conform to the scoring rules of the Peds FACIT Fatigue instrument. The scores in the second column correspond to the scores used in the full PROsetta Stone report.

8. Appendix Table 15: Indirect (Raw to Raw to Scale) Equipercentile Crosswalk Table – Pediatric FACIT Fatigue to PROMIS Pediatric Fatigue – Table 13 is recommended

| Peds FACIT Fatigue Actual Score* | Peds FACIT Fatigue Raw Score* | Equipercentile PROMIS Scaled Score Equivalents (No Smoothing) | Equipercentile Equivalents with Postsmoothing (Less Smoothing) | Equipercentile Equivalents with Postsmoothing (More Smoothing) |
|---|--------------------------------------|--|---|---|
| 52 | 13 | 27 | 26 | 23 |
| 51 | 14 | 30 | 30 | 30 |
| 50 | 15 | 33 | 33 | 33 |
| 49 | 16 | 36 | 36 | 36 |
| 48 | 17 | 39 | 38 | 39 |
| 47 | 18 | 41 | 41 | 41 |
| 46 | 19 | 43 | 43 | 43 |
| 45 | 20 | 45 | 45 | 45 |
| 44 | 21 | 47 | 48 | 47 |
| 43 | 22 | 50 | 49 | 49 |
| 42 | 23 | 50 | 50 | 50 |
| 41 | 24 | 51 | 51 | 51 |
| 40 | 25 | 52 | 52 | 52 |
| 39 | 26 | 53 | 53 | 53 |
| 38 | 27 | 54 | 54 | 54 |
| 37 | 28 | 55 | 55 | 55 |
| 36 | 29 | 56 | 56 | 56 |
| 35 | 30 | 58 | 57 | 57 |
| 34 | 31 | 58 | 58 | 58 |
| 33 | 32 | 58 | 58 | 58 |
| 32 | 33 | 59 | 59 | 59 |
| 31 | 34 | 60 | 60 | 60 |
| 30 | 35 | 60 | 60 | 60 |
| 29 | 36 | 61 | 61 | 61 |
| 28 | 37 | 61 | 61 | 62 |
| 27 | 38 | 62 | 62 | 62 |
| 26 | 39 | 63 | 63 | 63 |
| 25 | 40 | 64 | 64 | 64 |
| 24 | 41 | 66 | 65 | 65 |
| 23 | 42 | 66 | 66 | 66 |
| 22 | 43 | 67 | 67 | 66 |
| 21 | 44 | 67 | 68 | 67 |
| 20 | 45 | 68 | 68 | 68 |
| 19 | 46 | 69 | 69 | 69 |
| 18 | 47 | 70 | 70 | 70 |
| 17 | 48 | 71 | 70 | 71 |
| 16 | 49 | 72 | 71 | 72 |
| 15 | 50 | 72 | 72 | 72 |
| 14 | 51 | 73 | 73 | 73 |
| 13 | 52 | 73 | 74 | 74 |

PROsetta Stone® - PROMIS Pediatric Fatigue and Pediatric FACIT Fatigue

| | | | | |
|----|----|----|----|----|
| 12 | 53 | 75 | 75 | 75 |
| 11 | 54 | 77 | 77 | 76 |
| 10 | 55 | 78 | 78 | 78 |
| 9 | 56 | 81 | 80 | 79 |
| 8 | 57 | 86 | 81 | 79 |
| 7 | 58 | 86 | 81 | 80 |
| 6 | 59 | 86 | 82 | 81 |
| 5 | 60 | 86 | 83 | 82 |
| 4 | 61 | 86 | 83 | 83 |
| 3 | 62 | 86 | 84 | 84 |
| 2 | 63 | 86 | 85 | 84 |
| 1 | 64 | 86 | 85 | 85 |
| 0 | 65 | 86 | 86 | 86 |

*The scores in the first column conform to the scoring rules of the Peds FACIT Fatigue instrument. The scores in the second column correspond to the scores used in the full PROsetta Stone report.